

**Loewner – type Parametric Representation
of univalent self-maps with
given boundary regular fixed points**

PAVEL GUMENYUK

UNIVERSITY OF STAVANGER, NORWAY

**Tokyo Institute of Technology
Japan, November 23, 2014**

One of the most classical object of study
in *Geometric Function Theory* is the

Class \mathcal{S}

By \mathcal{S} we denote the class of all *holomorphic functions*

$$f : \mathbb{D} \xrightarrow{\text{into}} \mathbb{C}, \quad \mathbb{D} := \{z : |z| < 1\},$$

which are

- ▶ *univalent* in \mathbb{D} , and
- ▶ normalized by the condition $f(0) = 0, f'(0) = 1$.

The study of \mathcal{S} is difficult in many aspects, in particular, because:

- there is **no** natural **linear structure** in the class \mathcal{S} ;
- the class \mathcal{S} is even **not** a **convex** set in $\text{Hol}(\mathbb{D}, \mathbb{C})$.

Theorem (Parametric Representation) [Loewner, 1923;
Kufarev, 1943; Pommerenke, 1965-75; Gutlyanski, 1970]

$$\mathcal{S} = \left\{ \mathbb{D} \ni z \mapsto f[p](z) = \lim_{t \rightarrow +\infty} e^t \varphi_{0,t}(z) : \right. \\ \left. [0, +\infty) \ni t \mapsto \varphi_{0,t}(z) =: w(t) \text{ solves (1)} \right\}$$

(classical radial) Loewner–Kufarev ODE

$$\frac{dw}{dt} = -w(t)p(w(t), t), \quad w(0) = z \in \mathbb{D}, \quad (1)$$

where $p : \mathbb{D} \times [0, +\infty) \rightarrow \mathbb{C}$ is a *classical Herglotz function*, i.e.

- ▶ $p(z, \cdot)$ is measurable for all $z \in \mathbb{D}$;
- ▶ $p(\cdot, t)$ is holomorphic for all $t \geq 0$;
- ▶ $\operatorname{Re} p > 0$ and $p(0, t) = 1$ for all $t \geq 0$.

Thus the Parametric Representation is the *surjective map* $p \mapsto f[p]$ from the *convex cone* \mathcal{P}_0 of all classical Herglotz functions onto \mathcal{S} .

Notation: $\mathcal{U}(\mathbb{D}) := \{\varphi \in \text{Hol}(\mathbb{D}, \mathbb{D}) : \varphi \text{ is univalent}\}$,
 $\mathcal{U}_0(\mathbb{D}) := \{\varphi \in \mathcal{U}(\mathbb{D}) : \varphi(0) = 0, \varphi'(0) > 0\}$ — **semigroups!**

Theorem (“Parametric Representation folklore”)

Let $\varphi : \mathbb{D} \rightarrow \mathbb{C}$. Then $\varphi \in \mathcal{U}_0(\mathbb{D}) \iff \boxed{\varphi = \varphi_{0,T}}$,
where $T := -\log \varphi'(0)$ and $t \mapsto \varphi_{0,t}(z) =: w(t)$ is the solution to

$$dw(t)/dt = -w(t)p(w(t), t), \quad w(0) = z \in \mathbb{D},$$

for some classical Herglotz function p .

Simple idea to represent $\mathcal{U}(\mathbb{D})$

For $\varphi \in \mathcal{U}(\mathbb{D})$, write $\varphi = L \circ \varphi_0$, where $\varphi_0 \in \mathcal{U}_0(\mathbb{D})$ and $L \in \text{Möb}(\mathbb{D})$.

Not every simple idea turns out to be productive.

Decomposition $\varphi = L \circ \varphi_0$ does not allow to study infinitesimal structure of subsemigroups in $\mathcal{U}(\mathbb{D})$ and $\text{Hol}(\mathbb{D}, \mathbb{D})$.

Examples of semigroups w.r.t. $\cdot \circ \cdot$.

✎ Probability generating functions of Galton – Watson processes

$$\mathcal{H}_{\text{gen}} := \left\{ \varphi(z) = \sum_{n=0}^{+\infty} p_n z^n : p_n \geq 0, \sum_{n=0}^{+\infty} p_n = 1 \right\};$$

✎ $\mathcal{H}_{\infty}(\mathbb{H}) := \{ \varphi \in \text{Hol}(\mathbb{H}, \mathbb{H}) : \text{Im } \varphi(z) \geq \text{Im } z \}$, $\mathbb{H} := \{z : \text{Im } z > 0\}$,

$$\mathcal{H}_{\infty}(\mathbb{H}) = \{\text{id}_{\mathbb{H}}\} \cup \left\{ \varphi \in \text{Hol}(\mathbb{H}, \mathbb{H}) : \varphi^{\circ n} \xrightarrow{n \rightarrow \infty} \infty \right\};$$

✎ The reciprocal Cauchy transforms of probabil. measures with

$$\text{finite variance and mean zero } \mathcal{H}_{\text{CT}}(\mathbb{H}) := \left\{ \varphi(z) = \left(\int_{\mathbb{R}} \frac{d\mu(x)}{z-x} \right)^{-1} :$$

$$\mu \geq 0, \mu(\mathbb{R}) = 1, \int_{\mathbb{R}} x^2 d\mu(x) < +\infty, \int_{\mathbb{R}} x d\mu(x) = 0 \right\} \subset \mathcal{H}_{\infty}(\mathbb{H});$$

Examples of semigroups — CONTINUED

☞ $\mathcal{H}_{\text{SLE}}(\mathbb{H}) := \{\varphi \in \text{Hol}(\mathbb{H}, \mathbb{H}) \text{ meromorphic at } \infty :$

$$\varphi(z) = z + \sum_{n=1}^{+\infty} c_n/z^n, c_n \in \mathbb{R}\} \subset \mathcal{H}_{\text{CT}}(\mathbb{H});$$

☞ $\mathcal{H}_{0,1}(\mathbb{D}) := \{\varphi \in \text{Hol}(\mathbb{D}, \mathbb{D}) \text{ with } \varphi(0) = 0 \text{ and BRFP at } 1\};$

☞ “Unival. analogues”: $\mathcal{U}_{\text{gen}}, \mathcal{U}_{\infty}(\mathbb{H}), \mathcal{U}_{\text{CT}}(\mathbb{H}), \mathcal{U}_{\text{SLE}}(\mathbb{H}), \mathcal{U}_{0,1}(\mathbb{D})$.

Definition (BRFP)

$\sigma \in \mathbb{T}$ is a *boundary regular fixed point (BRFP)* of $\varphi \in \text{Hol}(\mathbb{D}, \mathbb{D})$ if

$$\angle \lim_{z \rightarrow \sigma} \varphi(z) = \sigma, \quad \varphi'(\sigma) := \angle \lim_{z \rightarrow \sigma} \frac{\varphi(z) - \sigma}{z - \sigma} \neq \infty.$$

V. V. Goryainov, 1987, 1992, 1996, 2002(2), 2011: *infinitesimal structure* and *parametric representation* of semigroups in $\text{Hol}(\mathbb{D}, \mathbb{D})$

Definition (one-parameter semigroups)

A *one-parameter semigroup* $(\phi_t) \subset \text{Hol}(\mathbb{D}, \mathbb{D})$ is a continuous semigroup homomorphism

$$([0, +\infty), \cdot + \cdot) \ni t \mapsto \phi_t \in (\text{Hol}(\mathbb{D}, \mathbb{D}), \cdot \circ \cdot).$$

One-parameter semigroups as semiflows

For any one-parameter semigroup (ϕ_t) there exists $G \in \text{Hol}(\mathbb{D}, \mathbb{C})$ s.t.

$$\frac{d\phi_t(z)}{dt} = G(\phi_t(z)), \quad \phi_0(z) = z, \quad \text{for all } z \in \mathbb{D} \text{ and all } t \geq 0. \quad (4)$$

Definition: G is called the *(infinitesimal) generator* of (ϕ_t) .

Clearly,

$G \in \text{Hol}(\mathbb{D}, \mathbb{C})$ is an infinitesimal generator
(of some one-parameter semigroup)

$\iff G$ is a *semicomplete* vector field in \mathbb{D} , i.e. for any $z \in \mathbb{D}$,

the I.V.P. $dw(t)/dt = G(w(t)), \quad w(0) = z, \quad (5)$

has a unique solution $w = w(t)$ defined for all $t \in [0, +\infty)$.

Hence there is a *1-to-1 correspondence* between

- one-parameter semigroups (ϕ_t) , and
- semicomplete holomorphic vector fields $G : \mathbb{D} \rightarrow \mathbb{C}$.

Definition (Infinitesimal structure of a semigroup)

Let $U \subset \text{Hol}(\mathbb{D}, \mathbb{D})$ be a (sub)semigroup. The set $\mathcal{G}[U]$ of all infinitesimal generators of one-parameter semigroups $(\phi_t) \subset U$ is called the *infinitesimal structure* of the semigroup U .

Recall the notation:

$$\mathcal{U}(\mathbb{D}) := \{\varphi \in \text{Hol}(\mathbb{D}, \mathbb{D}) : \varphi \text{ univalent}\}$$

$$\mathcal{U}_0(\mathbb{D}) := \{\varphi \in \mathcal{U}(\mathbb{D}) : \varphi(0) = 0, \varphi'(0) > 0\}$$

$$\mathcal{U}_\infty(\mathbb{H}) := \{\varphi \in \text{Hol}(\mathbb{H}, \mathbb{H}) : \varphi \text{ univalent, } \text{Im } \varphi(z) \geq \text{Im } z\}$$

- ☞ All semicomplete holomorphic vector fields are given by the **Berkson – Porta** formula:

$$\mathcal{G} := \mathcal{G}[\text{Hol}(\mathbb{D}, \mathbb{D})] = \mathcal{G}[\mathcal{U}(\mathbb{D})] = \left\{ G(z) = (\tau - z)(1 - \bar{\tau}z)p(z) : \right. \\ \left. \tau \in \overline{\mathbb{D}}, p \in \text{Hol}(\mathbb{D}, \mathbb{C}), \text{Re } p \geq 0 \right\};$$

- ☞ $\mathcal{G}[\mathcal{U}_0(\mathbb{D})] = \left\{ G(z) = -zp(z) : p \in \text{Hol}(\mathbb{D}, \mathbb{C}), \text{Re } p \geq 0, \right. \\ \left. \text{Im } p(0) = 0 \right\};$

- ☞ $\mathcal{G}[\mathcal{U}_\infty(\mathbb{H})] = \mathcal{G}[\mathcal{H}_\infty(\mathbb{H})] = \left\{ G \in \text{Hol}(\mathbb{H}, \mathbb{C}) : \text{Im } G \geq 0 \right\};$

- ☞ $\mathcal{G}[\mathcal{U}_{\text{CT}}(\mathbb{H})] = \left\{ G(z) = \int_{\mathbb{R}} \frac{d\mu(x)}{x-z} : 0 \leq \mu < +\infty \text{ Borel measure} \right\}.$

Denote: (ϕ_t^G) the one-param. semigroup generated by $G \in \mathcal{G}$.

Analogue of the Lie exponential map

$$\mathcal{G}[U] \ni G \mapsto \text{Exp}_{\text{Lie}}(G) := \phi_1^G \in U \subset \text{Hol}(\mathbb{D}, \mathbb{D}) \text{ (subsemigroup)}$$

- 😊 For Lie groups, the **Exp**-map recovers the group (*at least locally*)
- 😞 However in our case, typically $\text{Exp}_{\text{Lie}}(\mathcal{G}[U]) \neq U, \neq \mathcal{O}_U(\text{id}_{\mathbb{D}})$.

Loewner's idea: Instead of (ϕ_t) 's satisfying the autonomous ODE

$$d\phi_t(z)/dt = G(\phi_t(z)), \quad t \geq 0, \quad \phi_0(t) = z \in \mathbb{D}, \quad (7)$$

consider two-parameter families $(\varphi_{s,t})_{t \geq s \geq 0}$, generated by its **non-autonomous analogue:**

$$d\varphi_{s,t}(z)/dt = G(\varphi_{s,t}(z), t), \quad t \geq s \geq 0, \quad \varphi_{s,s}(z) = z \in \mathbb{D}, \quad (8)$$

where $G(\cdot, t) \in \mathcal{G}[U]$ for a.e. $t \geq 0$

(plus a “**reasonable assumption**” regarding dependence on t).

- ▶ The ODE $\frac{d\varphi_{s,t}(z)}{dt} = G(\varphi_{s,t}(z), t)$, $t \geq s \geq 0$, $\varphi_{s,s}(z) = z$, (8) is called the *general Loewner equation*;
- ▶ the functions $G : [0, +\infty) \ni t \mapsto G(\cdot, t) \in \mathcal{G}$ are called *Herglotz vector fields*.
- ▶ the families $(\varphi_{s,t})$ are usually referred to as *evolution families*.

Definition

We say that a semigroup $U \subset \text{Hol}(\mathbb{D}, \mathbb{D})$ admits a *Loewner-type representation* if the union $\mathcal{R}[U]$ of all evolution families $(\varphi_{s,t})$ generated by Herglotz vector fields G with $G(\cdot, t) \in \mathcal{G}[U]$ for a.e. $t \geq 0$ coincides with U .

This means that one can reconstruct the semigroup U from its infinitesimal structure $\mathcal{G}[U]$ using the general Loewner ODE (8).
But, a priori, this depends on the *strict definition* of Herglotz v. f.'s.

Given:

- an *abstract* topological semigroup U , and
- an appropriate notion of *differentiability* and of the *derivative* for maps $\mathbb{R} \supset [a, b) \rightarrow U$,

one can consider the *analogue of the general Loewner ODE*
and hence construct the *Loewner – type representation* for U .

We assume, in particular, that:

► *Differentiability of re-parametrizations:*

- If $\psi: [a, b) \rightarrow U$ and $\tau: [c, d) \rightarrow [a, b)$ are differentiable, then $\psi \circ \tau: [c, d) \rightarrow U$ is also differentiable.
- If $\tau'(t_0) = 1$, for some $t_0 \in [c, d)$, then $(\psi \circ \tau)'(t_0) = \psi'(\tau(t_0))$.

► *Differentiability of the right translation $\mathcal{T}_\varphi: \psi \mapsto \psi\varphi$, $\varphi \in U$ fixed:*

- If $\psi: [a, b) \rightarrow U$ is differentiable, then so is $t \mapsto \mathcal{T}_\varphi\psi$.
- If $\psi_1(t_1) = \psi_2(t_2)$ and $\psi'_1(t_1) = \psi'_2(t_2)$, for some $t_1, t_2 \in [a, b)$, then $(\mathcal{T}_\varphi\psi_1)'(t_1) = (\mathcal{T}_\varphi\psi_2)'(t_2)$ [and, trivially, $\mathcal{T}_\varphi\psi_1(t_1) = \mathcal{T}_\varphi\psi_2(t_2)$].

In this abstract setting, the *infinitesimal structure* is

$$\mathcal{G}[U] := \left\{ \left. \frac{d}{dt} \phi_t \right|_{t=0} : (\phi_t) \subset U \text{ a differentiable 1-param. semigroup} \right\}.$$

Remark: if, e.g., U is a Lie group, then $\mathcal{G}[U] = T_{\text{id}} U$.

☞ The *differential of the right translation* can be defined as follows:

$$d_0 \mathcal{T}_\varphi \left(\left. \frac{d}{dt} \phi_t \right|_{t=0} \right) := \left. \frac{d}{dt} \mathcal{T}_\varphi \phi_t \right|_{t=0} = \left. \frac{d}{dt} \phi_t \varphi \right|_{t=0}.$$

Remark: for $U \subset \text{Hol}(\mathbb{D}, \mathbb{D})$, $\mathcal{G}[U] \subset \text{Hol}(\mathbb{D}, \mathbb{C})$ and $d_0 \mathcal{T}_\varphi(G) = G \circ \varphi$.

☞ Any differentiable 1-param. semigroup satisfies [by the very definition]

$$\left. \frac{d}{dt} \phi_t \right|_{t=0} = d_0 \mathcal{T}_{\phi_t}(G), \quad t \geq 0, \quad \phi_0 = \text{id}, \quad (9)$$

where $\mathcal{G}[U] \ni G := \left. \frac{d}{dt} \phi_t \right|_{t=0}$. The *analogue of the Loewner ODE*

would be $\left. \frac{d}{dt} \varphi_{s,t} \right|_{t=0} = d_0 \mathcal{T}_{\varphi_{s,t}}(G(t)), \quad t \geq s \geq 0, \quad \varphi_{s,s} = \text{id}, \quad (10)$

where $G : [0, +\infty) \rightarrow \mathcal{G}[U]$.

Definition

The semigroup U is said to *admit a Loewner – type representation*, if there exists a class H of functions $G : [0, +\infty) \rightarrow \mathcal{G}[U]$ s.t.:

- (i) given any $G \in H$, for every $s \geq 0$ the I.V.P. the Loewner ODE

$$\frac{d}{dt}\varphi_{s,t} = d_0\mathcal{T}_{\varphi_{s,t}}(G(t)), \quad \varphi_{s,s} = \text{id}, \quad (11)$$

has a unique solution $t \mapsto \varphi_{s,t}^G$ defined for all $t \geq s$;

- (ii) for any $\phi \in U$ there exists $G \in H$ such that $(\varphi_{s,t}^G) \ni \phi$.

☞ Ch. Loewner himself applied this scheme and obtained a Loewner – type representation for a certain semigroup of matrices.

[*On totally positive matrices*, Math. Z. **63** (1955), 338–340]

- ✓ In that case, of course, *no trouble with differentiability* appears, because the semigroup \subset a finite-dimensional linear space.
- ✓ Nevertheless, this shows that

this abstract scheme can work in different settings.

Keeping in mind these abstract ideas, let us return to $U \subset \text{Hol}(\mathbb{D}, \mathbb{D}) \dots$

- ✓ V.V. Goryainov, for several *concrete choices* of the semigroup $U \subset \text{Hol}(\mathbb{D}, \mathbb{D})$, gave precise definitions of Herglotz vector fields, *specific for each U* , and established a Loewner – type representation in each case.
- ✓ Recently, another *general approach* has been suggested by F. Bracci, M.D. Contreras and S. Díaz-Madrigal, 2008
[*J. Reine Angew. Math.* **672**(2012), 1–37]

Definition (Bracci *et al*) [for the whole semigroup $\text{Hol}(\mathbb{D}, \mathbb{D})$]

A function $G : \mathbb{D} \times [0, +\infty) \rightarrow \mathbb{C}$ is a *Herglotz vector field* if:

- (i) $G(\cdot, t) \in \mathcal{G}$ for a.e. $t \geq 0$;
- (ii) $G(z, \cdot)$ is measurable on $[0, +\infty)$ for every $z \in \mathbb{D}$;
- (iii) for any compact set $K \subset \mathbb{D}$,
 $M_K(t) := \sup_K |G(\cdot, t)|$ is locally integrable on $[0, +\infty)$.

Definition (evolution families — *intrinsic definition*) [Bracci et al]

A family $(\varphi_{s,t})_{t \geq s \geq 0} \subset \text{Hol}(\mathbb{D}, \mathbb{D})$ is called an *evolution family* if:

- (i) $\varphi_{s,s} = \text{id}_{\mathbb{D}}$ for all $s \geq 0$;
- (ii) $\varphi_{s,t} = \varphi_{u,t} \circ \varphi_{s,u}$ whenever $t \geq u \geq s \geq 0$;
- (iii) for any $z \in \mathbb{D}$, the maps $[s, +\infty) \ni t \mapsto \varphi_{s,t}(z)$ are locally absolutely continuous *uniformly w.r.t.* $s \geq 0$.

Theorem (Bracci et al)

The general Loewner ODE

$$d\varphi_{s,t}(z)/dt = G(\varphi_{s,t}(z), t), \quad t \geq s \geq 0, \quad \varphi_{s,s}(z) = z, \quad (8)$$

establishes an (essentially) *1-to-1 correspondence* between *Herglotz vector fields* G and *evolution families* $(\varphi_{s,t})$.

This includes uniqueness and global existence for solutions to (8).

Note: (8) is to be understood as a *Carathéodory ODE*.

Problem: *construct a Loewner – type parametric representation for semigroups formed by univalent self-maps with given fixed points.*

Let \mathcal{F} be a finite set of points on $\mathbb{T} := \partial\mathbb{D}$.

First family of semigroups

$$\mathcal{U}(\mathbb{D}, \mathcal{F}) := \left\{ \varphi \in \mathcal{U}(\mathbb{D}) : \text{each } \sigma \in \mathcal{F} \text{ is a BRFP of } \varphi \right\}$$

“BRFP” = “*boundary regular fixed point*”:

A point $\sigma \in \partial\mathbb{D}$ is said to be *BRFP* of $\varphi \in \text{Hol}(\mathbb{D}, \mathbb{D})$ if

$$\exists \angle \lim_{z \rightarrow \sigma} \varphi(z) = \sigma \quad \text{and} \quad \exists \varphi'(\sigma) := \angle \lim_{z \rightarrow \sigma} \frac{\varphi(z) - \sigma}{z - \sigma} \neq \infty.$$

Presence of a BRFPs affects the behaviour of the map at the internal points $z \in \mathbb{D}$. The basic and most classical result is

Theorem (Julia – Wolff – Carathéodory)

Suppose that $\varphi \in \text{Hol}(\mathbb{D}, \mathbb{D})$, $\sigma \in \partial\mathbb{D}$ and $\exists \angle \lim_{z \rightarrow \sigma} \varphi(z) = \sigma$.

Then **TFAE**:

- (i) σ is a BRFP of φ ;
- (ii) $\alpha_\varphi(\sigma) := \liminf_{z \rightarrow \sigma} \frac{1 - |\varphi(z)|}{1 - |z|} < +\infty$;
- (iii) there exists $A < +\infty$ such that

$$\frac{|\varphi(z) - \sigma|^2}{1 - |\varphi(z)|^2} \leq A \frac{|z - \sigma|^2}{1 - |z|^2} \quad \text{for all } z \in \mathbb{D}. \quad \text{[Julia's inequality]}$$

Moreover, the minimal value of A ,
the *boundary dilation coefficient* $\alpha_\varphi(\sigma)$ and $\varphi'(\sigma)$ all coincide.

Fix additionally $\tau \in \overline{\mathbb{D}} \setminus \mathcal{F}$.

Second family of semigroups

$$\mathcal{U}_\tau(\mathbb{D}, \mathcal{F}) := \{\text{id}_{\mathbb{D}}\} \cup \{\varphi \in \mathcal{U}(\mathbb{D}, \mathcal{F}) \setminus \{\text{id}_{\mathbb{D}}\} : \tau \text{ is the DW-point of } \varphi\}$$

“DW-point” = “*Denjoy–Wolff point*” [Denjoy–Wolff Theorem]

For any $\varphi \in \text{Hol}(\mathbb{D}, \mathbb{D}) \setminus \{\text{id}_{\mathbb{D}}\}$,

$\exists!$ (boundary regular) fixed point $\tau \in \overline{\mathbb{D}}$ such that $|\varphi'(\tau)| \leq 1$.

Moreover, if φ is *not* an elliptic automorphism of \mathbb{D} ,

then $\varphi^{on} \rightarrow \tau$ l.u. in \mathbb{D} as $n \rightarrow +\infty$.

This point τ is called the *Denjoy–Wolff point* of φ .

- ☞ Loenwer's idea potentially can work in the general setting of an abstract semigroup with “compatible diffeology”.
However, no criteria for such a semigroup to admit a parametric representation is known.
So it is interesting to study more examples.
- ☞ In Geometric Function Theory there has been considerable interest to study self-maps with given BRFP's
H. Unkelbach, 1938, 1940; C. Cowen, Ch. Pommerenke, 1982;
Ch. Pommerenke, A. Vasil'ev, 2000; J.M. Anderson, A. Vasil'ev, 2008;
M. Elin, D. Shoikhet, N. Tarkhanov, 2011;
V.V. Goryainov [talk at Steklov Math. Inst., Moscow, 26/12/2011];
A. Frolova, M. Levenshtein, D. Shoikhet, A. Vasil'ev, ArXiv:1309.3074, 2013.
- ☞ The infinitesimal structure of $\mathcal{U}(\mathbb{D}, \mathcal{F})$ and $\mathcal{U}_\tau(\mathbb{D}, \mathcal{F})$ is well-studied.
- ☞ The following result: [see next slide]

Theorem (Bracci, Contreras, Díaz-Madrugal, **P. Gum.**, 2013)

Let $(\varphi_{s,t})$ be an evolution family with Herglotz vector field G .

Let $\sigma \in \mathbb{T}$. Then **TFAE**:

(i) σ is a BRFP of $\varphi_{s,t}$ for all $t \geq s \geq 0$;

(ii) G satisfies:

(ii.1) for a.e. $t \geq 0$ fixed, $G(\cdot, t) \in \mathcal{G}[\mathcal{U}(\mathbb{D}, \{\sigma\})]$ [Contreras, Díaz-Madrugal, Pommerenke, 2006] $\iff \exists \lambda \lim_{z \rightarrow \sigma} \frac{G(t, z)}{z - \sigma} =: \lambda(t) \neq \infty$;

(ii.2) $\lambda(t)$ is locally integrable on $[0, +\infty)$.

If the above conditions hold, then $\varphi'_{s,t}(\sigma) = \exp(\int_s^t \lambda(\xi) d\xi)$.

Let U be our semigroup, $\mathcal{U}(\mathbb{D}, \mathcal{F})$ or $\mathcal{U}_\tau(\mathbb{D}, \mathcal{F})$.

This theorem **reduces our problem to checking** whether

$$\mathcal{R}[U] := \bigcup \{\varphi_{s,t}\} \stackrel{?}{=} U,$$

where the **union** is taken over **all evolution families** $\{\varphi_{s,t}\} \subset U$.

Theorem (P. Gum. — work in progress)

Let $\mathcal{F} \subset \mathbb{T}$ be a finite set, $n := \text{Card}(\mathcal{F})$, and $\tau \in \overline{\mathbb{D}}$.

The following semigroups U admit

the Loewner-type parametric representation, i.e. $\mathcal{R}[U] = U$:

- ✓ $U = \mathcal{U}_\tau(\mathbb{D}, \mathcal{F})$ for $\tau \in \mathbb{D}$ and $n = 1$; [Unkelbach and Goryainov]
- ✓ $U = \mathcal{U}_\tau(\mathbb{D}, \mathcal{F})$ for $\tau \in \mathbb{T}$ and $n \leq 2$;
- ✓ $U = \mathcal{U}(\mathbb{D}, \mathcal{F})$ for $n \leq 3$.

H. Unkelbach, 1940: an attempt to give

the Loewner-type parametric representation for $\mathcal{U}_0(\mathbb{D}, \{1\})$;

V.V. Goryainov, approx. 2013 (to appear in *Mat. Sb.*):

the complete proofs.

Conjecture [know how to prove]

If $\tau \in \mathbb{D}$, then the semigroup $\mathcal{U}_\tau(\mathbb{D}, \mathcal{F})$ admits
the Loewner type representation for any finite set $\mathcal{F} \subset \mathbb{T}$.

Open problem

Given a finite $\mathcal{F} \subset \mathbb{T}$ with $\text{Card}(\mathcal{F}) = n$,

- ❓ Does the semigroups $\mathcal{U}_\tau(\mathbb{D}, \mathcal{F})$ admits
the Loewner type representation for $\tau \in \mathbb{T}$ and $n > 2$?
- ❓ Does the semigroups $\mathcal{U}(\mathbb{D}, \mathcal{F})$ admits
the Loewner type representation for $n > 3$?

My conjecture is that the correct answer for both questions is NO.

有難うございます。

ARIGATOUGOZAIMASU